

Baryogenesis from dark matter in an inflationary universe

Wan-Zhe Feng,¹ Anupam Mazumdar,² and Pran Nath³

¹*Department of Physics and Institute for Advanced Study,
The Hong Kong University of Science and Technology, Hong Kong*

²*Consortium for fundamental physics, Lancaster University, LA1 4YB, United Kingdom*

³*Department of Physics, Northeastern University, Boston, MA 02115, USA*

(Dated: February 4, 2013)

We consider the possibility that in an inflationary universe, the inflaton field decays purely into the dark sector creating asymmetric dark matter at the end of inflation. This asymmetry is subsequently transmuted into leptons and baryons. We consider this possibility in the framework of a generic inflation model, and compute the amount of asymmetric dark matter created from the out of equilibrium decays of the inflaton with CP violating Yukawa couplings. The dark matter asymmetry is then transferred to the visible sector by the asymmetry transfer equation and generates an excess of $B - L$. Baryogenesis occurs via sphaleron processes which conserve $B - L$ but violate $B + L$. A mechanism for the annihilation of the symmetric component of dark matter is also discussed. The model leads to multi-component dark matter consisting of both bosonic and fermionic components.

Keywords: Asymmetric Dark Matter, Inflation, Leptogenesis, Baryogenesis

PACS numbers: 95.35.+d, 12.60.Jv

Introduction: In this work we discuss the possibility that in an inflationary universe [1, 2] the inflaton field decays into the dark sector and creates an asymmetry, instead of decaying into standard model particles. The dark matter asymmetry then transmutes into leptons and baryons. As is well known the generation of a baryon excess (or a lepton excess) requires satisfaction of the three Sakharov conditions [3]: a violation of baryon (or lepton) number, existence of C and CP violation, and non-equilibrium processes. Baryon and lepton number violations appear in the Standard Models and its supersymmetric extensions via higher dimensional operators (for a review see [4]) and they can also arise via spontaneous breaking [5]. Thus in the analysis of the genesis of dark matter via the decay of the inflaton field, we will assume the existence of such violations. The remaining Sakharov conditions are also met leading to the generation of asymmetric dark matter. A part of dark matter then transmutes to the visible sector. An analysis in similar spirit where dark matter is the genesis of visible matter was discussed in [6]. Our model is significantly different from this work and further we also discuss the genesis of asymmetric dark matter as arising from the decay of the inflaton field. The symmetric component of dark matter is depleted by mechanisms similar to those discussed in [7].

The model: We will work in a supersymmetric framework where the superpotential of the model is given by

$$W = W_{\text{inf}} + W_{\text{gen}} + W_{\text{tran}} + W_{\text{MSSM}}. \quad (1)$$

Here W_{inf} enters in the inflationary period, W_{gen} generates the asymmetry for the dark matter through the decay of the inflaton, W_{tran} transfers the asymmetry

from the dark sector to the visible sector, and W_{MSSM} is the superpotential of the minimal supersymmetric standard model. Regarding inflation, currently one does not have a standard model of inflation but there are several possible candidate scenarios [30]. To be concrete we will consider a simple possibility of slow roll inflation driven by D-terms [9, 10] but we do not go into details. However, we emphasize that our analysis is not dependent on the details of the inflationary model in any significant way as we can accommodate a variety of current inflationary models (for a review see [11]).

Genesis of asymmetric dark matter: We assume there exist several $\hat{N}_i$ fields ($i \geq 2$) in the early universe with masses $M_{\hat{N}_i}$, where $\hat{N} = (N, \tilde{N})$ and N is the Majorana field and $\tilde{N}$ is the super-partner field. The scalar field of the lightest $\hat{N}_i$ superfields plays the role of the inflaton, i.e., $\phi \equiv \tilde{N}_1$. We now focus on the period when inflation ends. Here the dynamics is driven by the coupling of the inflaton superfield $\hat{\phi}$ to the dark sector. The dark sector is comprised of $(\hat{X}, \hat{X}^c, \hat{X}', \hat{X}'^c)$ which are charged under the gauge group $U(1)_x$ with charges $(+1, -1, -1, +1)$. All of the MSSM fields are not charged under this new gauge symmetry $U(1)_x$. We assume the $\hat{N}_i$ superfields carry a non-vanishing lepton number $+2$, $\hat{X}, \hat{X}'$ carry lepton number -1 and $\hat{X}^c, \hat{X}'^c$ carry lepton number $+1$. W_{gen} is invariant under both $U(1)_x$ and lepton number [31] and it takes the following form

$$W_{\text{gen}} = \lambda_i \hat{N}_i \hat{X} \hat{X}' + m \hat{X} \hat{X}^c + m' \hat{X}' \hat{X}'^c, \quad (2)$$

where the Yukawa coupling λ_i is assumed to be complex. The interaction of W_{gen} describes the genesis of




FIG. 1: Loop diagrams which are responsible for the genesis of asymmetric dark matter from the decay of N_i to final states $X\bar{X}'$ and there are similar diagrams for the decay of the N_i to the final states $\bar{X}X'$, and for the decay of $\tilde{N}_i$ to XX' and to $\bar{X}\bar{X}'$.

dark matter. It gives rise to the decays

$$N_i \rightarrow X\bar{X}', \bar{X}X', \bar{X}\bar{X}'^*, \bar{X}^*\bar{X}', \quad \tilde{N}_i \rightarrow XX', \bar{X}\bar{X}'. \quad (3)$$

In the simplest model we have $i = 2$, and we assume $\tilde{N}_2$ mass M_2 is much larger than $\tilde{N}_1$ mass M_1 , so the generation of the asymmetry in the dark sector is mostly through $\tilde{N}_1$.

The genesis of the dark matter asymmetry arises from the interference of the one-loop diagrams shown in Fig. 1 with the tree-level diagrams similar to the conventional leptogenesis diagrams [12–15]. The asymmetries, i.e., the excess of $\hat{X}, \hat{X}'$ over their anti-particles $\bar{\hat{X}}, \bar{\hat{X}'}$ are measured by

$$\epsilon_{X\bar{X}'} = \frac{\Gamma(N_1 \rightarrow X\bar{X}') - \Gamma(N_1 \rightarrow \bar{X}\bar{X}'^*)}{\Gamma(N_1 \rightarrow X\bar{X}') + \Gamma(N_1 \rightarrow \bar{X}\bar{X}'^*)}, \quad (4)$$

$$\epsilon_{\bar{X}X'} = \frac{\Gamma(N_1 \rightarrow \bar{X}X') - \Gamma(N_1 \rightarrow \bar{X}^*\bar{X}')}{\Gamma(N_1 \rightarrow \bar{X}X') + \Gamma(N_1 \rightarrow \bar{X}^*\bar{X}')}, \quad (5)$$

$$\epsilon_{XX'} = \frac{\Gamma(\tilde{N}_1 \rightarrow XX') - \Gamma(\tilde{N}_1 \rightarrow \bar{X}\bar{X}')}{\Gamma(\tilde{N}_1 \rightarrow XX') + \Gamma(\tilde{N}_1 \rightarrow \bar{X}\bar{X}')}, \quad (6)$$

$$\epsilon_{\bar{X}\bar{X}'} = \frac{\Gamma(\tilde{N}_1 \rightarrow \bar{X}\bar{X}') - \Gamma(\tilde{N}_1 \rightarrow \bar{X}^*\bar{X}'^*)}{\Gamma(\tilde{N}_1 \rightarrow \bar{X}\bar{X}') + \Gamma(\tilde{N}_1 \rightarrow \bar{X}^*\bar{X}'^*)}. \quad (7)$$

There are two types of loops involved: vertex contribution and wave contribution as shown in Fig 1. It's straight forward to compute the asymmetry parameters $\epsilon_{X\bar{X}'}$, etc defined above. It turns out the contributions of the vertex diagrams and the wave diagrams satisfy the following relations [32]

$$\epsilon_{X\bar{X}'}^{vertex} = \epsilon_{\bar{X}X'}^{vertex} = \epsilon_{XX'}^{vertex} = \epsilon_{\bar{X}\bar{X}'}^{vertex} \equiv \epsilon^{vertex}, \quad (8)$$

$$\epsilon_{X\bar{X}'}^{wave} = \epsilon_{\bar{X}X'}^{wave} = \epsilon_{XX'}^{wave} = \epsilon_{\bar{X}\bar{X}'}^{wave} \equiv \epsilon^{wave}. \quad (9)$$

Specifically, we have [33]

$$\epsilon^{vertex} = -\frac{1}{32\pi} \frac{\text{Im}(\lambda_1^2 \lambda_2^{*2})}{|\lambda_1|^2} \frac{M_2}{M_1} \ln \frac{M_1^2 + M_2^2}{M_2^2}, \quad (10)$$

$$\epsilon^{wave} = -\frac{1}{32\pi} \frac{\text{Im}(\lambda_1^2 \lambda_2^{*2})}{|\lambda_1|^2} \frac{M_1(M_1 + M_2)}{M_2^2 - M_1^2}. \quad (11)$$

Thus the total asymmetry parameter is the sum of the vertex and the wave contributions as given by Eqs. (10), (11) and one has the following equalities

$$\epsilon_{X\bar{X}'} = \epsilon_{\bar{X}X'} = \epsilon_{XX'} = \epsilon_{\bar{X}\bar{X}'} \equiv \epsilon. \quad (12)$$

where ϵ is the sum of ϵ^{vertex} and ϵ^{wave} and in the limit $M_2 \gg M_1$, we obtain

$$\epsilon = \epsilon^{vertex} + \epsilon^{wave} \approx -\frac{1}{16\pi} \frac{\text{Im}(\lambda_1^2 \lambda_2^{*2})}{|\lambda_1|^2} \frac{M_1}{M_2}. \quad (13)$$

Thus the total excess of $X, \bar{X}, X', \bar{X}'$ over $\bar{X}, \bar{X}^*, \bar{X}', \bar{X}'^*$ generated by the decay of $\tilde{N}_1$ is given by:

$$\Delta n_X \equiv (n_{\hat{X}} - n_{\bar{\hat{X}}}) + (n_{\hat{X}'} - n_{\bar{\hat{X}'}}),$$

where Δn_X is computed to be

$$\Delta n_X = \left[\frac{3}{4} (\epsilon_{X\bar{X}'} + \epsilon_{\bar{X}X'}) + (\epsilon_{XX'} + \epsilon_{\bar{X}\bar{X}'}) \right] \frac{\kappa \zeta(3) g_N T^3}{\pi^2},$$

where $\zeta(3) \sim 1.202$. We may further write $\Delta n_X \approx 2\kappa s \epsilon / g_*$, where s is the entropy. Here the factor of $\frac{3}{4}$ is for N_i vs a factor of 1 for $\tilde{N}_i$, $g_N = 2$ are the degrees of freedom of the Majorana field, g_* is the entropy degrees of freedom for MSSM where $g_* \approx 228.75$, and κ is a washout factor due to inverse processes $X + \bar{X}', \bar{X} + X' \rightarrow N$ and $X + X', \bar{X} + \bar{X}' \rightarrow \tilde{N}$. A computation of κ requires solving the Boltzman equations [18]. The excess of $\hat{X}, \hat{X}'$ then give rise to a non-vanishing $(B - L)$ -number in the early universe:

$$(B - L)_t = (+1) \times \Delta n_X = 2\kappa s \epsilon / g_*, \quad (14)$$

where $(B - L)_t$ is the total $B - L$ in the Universe.

Leptogenesis and Baryogenesis from dark matter: We consider now supersymmetric interactions which can transfer a $B - L$ asymmetry from the dark sector to the visible sector, which give rise to both leptogenesis and baryogenesis. Leptogenesis is accomplished via a transfer equation similar to the one adopted in previous works [19]

$$W_{\text{tran}} = \frac{1}{M_{\text{asy}}^n} \hat{X} \hat{X}' \mathcal{O}_{\text{asy}}^{\text{MSSM}}. \quad (15)$$

As a specific example we consider the case

$$W_{\text{tran}} = \frac{1}{M_{\text{asy}}^2} \hat{X} \hat{X}' (L H_u)^2. \quad (16)$$

We assume that the transfer interaction is active only above the temperature T_{int} which lies above T_{SUSY} which corresponds to the SUSY breaking scale, i.e., $T_{\text{int}} > T_{\text{SUSY}} \sim \mathcal{O}(1\text{TeV})$. Of course there are other possible choices for T_{int} where W_{tran} would decouple (see [7] for a comprehensive discussion). To determine the generation of lepton number and baryon number from the couplings

of the dark sector to the visible sector we use the standard thermal equilibrium method introduced in [20]. At the scale $T_{\text{int}} > T_{\text{SUSY}}$ all the MSSM fields are ultra-relativistic and have non-vanishing chemical potential while the gauge bosons have a vanishing chemical potential. Constraints on the chemical potential arise from a variety of sources. The superpotential in MSSM reads

$$W_{\text{MSSM}} = g_u Q H_u U^c - g_d Q H_d D^c - g_e L H_d E^c + \mu H_u H_d \quad (17)$$

where the Yukawa sector gives the following constraints on the chemical potentials

$$\mu_{H_d} = -\mu_L - \mu_{E^c} = -\mu_Q - \mu_{D^c}, \quad (18)$$

$$\mu_{H_u} = -\mu_Q - \mu_{U^c}, \quad (19)$$

while the Higgs mixing term $\mu H_u H_d$ gives

$$\mu_{H_u} + \mu_{H_d} = 0. \quad (20)$$

Additionally, the sphaleron processes ($\prod Q_i Q_i Q_i L_i$, $i = 1, 2, 3$) give us the constraint,

$$3\mu_Q + \mu_L = 0, \quad (21)$$

and the condition that the total hypercharge of the Universe is zero leads to

$$Y = 3 \times (3\mu_Q - 6\mu_{U^c} + 3\mu_{D^c} - 3\mu_L + 3\mu_{E^c} + 2\mu_{H_u}) = 0. \quad (22)$$

Solving above equations, we can express all the chemical potentials in terms of the chemical potential of one single field, i.e., μ_L . The solutions show a generation of lepton and baryon number in the visible sector and one has

$$(B - L)_v = -\frac{237}{7}\mu_L, \quad (23)$$

where $(B - L)_v$ is the $B - L$ in the visible sector.

The dark matter also undergoes a readjustment as a consequence of thermal equilibrium and the residual dark matter after the action of the asymmetry transfer interaction Eq. (16) can be gotten via the chemical potential equation:

$$\mu_{\hat{X}} + \mu_{\hat{X}'} + 2(\mu_L + \mu_{H_u}) = 0. \quad (24)$$

Also the mass terms of $\hat{X}$, $\hat{X}^c$, $\hat{X}'$, $\hat{X}'^c$ in Eq. (2) at equilibrium give

$$\mu_{\hat{X}} + \mu_{\hat{X}^c} = \mu_{\hat{X}'} + \mu_{\hat{X}'^c} = 0, \quad (25)$$

which lead to

$$\mu_{\hat{X}} + \mu_{\hat{X}'} = -\mu_{\hat{X}^c} - \mu_{\hat{X}'^c} = -\frac{22}{7}\mu_L. \quad (26)$$

From the above we find

$$\sum_i X_i = -\frac{3 \times 2 \times 22}{7}\mu_L = \frac{44}{79}(B - L)_v. \quad (27)$$

where X_i are the number of dark matter particles of species i (in this case i takes on values from 1 – 4 since we have 4 species: $\hat{X}, \hat{X}^c, \hat{X}', \hat{X}'^c$). In Eq. (27) the factor of 3 = 1 + 2 counts the chemical potentials for both fermions and bosons of the superfields (from converting the μ 's to the excess of the number densities), and the factor of 2 counts the contribution also from $\hat{X}^c, \hat{X}'^c$. Using Eq. (27) it is also easy to find the ratio that

$$\frac{(B - L)_v}{(B - L)_t} \approx 0.64. \quad (28)$$

We can now determine the dark matter mass from the ratio of the dark matter relic density to baryonic relic density. Thus the ratio of dark matter relic density to the baryonic matter density is given by

$$\frac{\Omega_{\text{DM}}}{\Omega_{\text{B}}} = \frac{\sum_i X_i \cdot m_{\text{DM}}^i}{B \cdot m_{\text{B}}} \approx 5, \quad (29)$$

where m_{DM}^i are the masses of the dark matter particles and $m_{\text{B}} \sim 1$ GeV. There is an important subtlety here that although the total dark particle number is fixed after the asymmetry transfer interaction decouples, the total baryon number, however, keeps changing because of the sphaleron processes. As was explained in [7], the total baryon number to be used in this formula is B_{final} after the sphaleron processes decouple. Using Eq. (29) we have

$$m_{\text{DM}} = 5 \cdot \frac{B_{\text{final}}}{\sum_i X_i}, \quad (30)$$

where

$$B_{\text{final}} = \frac{30}{97}(B - L)_v \approx 0.31(B - L)_v. \quad (31)$$

Thus we obtain

$$m_{\text{DM}} \approx 2.78 \text{ GeV}. \quad (32)$$

From astrophysical constraints one has [21]

$$B_{\text{final}}/s \sim 6 \times 10^{-10}. \quad (33)$$

Using Eqs. (14), (28) and (31), we obtain

$$\epsilon \sim 4 \times 10^{-6}, \quad (34)$$

which sets bounds for complex couplings λ_i and the ratio M_1/M_2 . This can be seen by noting that Eq. (13) can be written in the form

$$\epsilon \approx -\frac{M_1}{16\pi M_2} |\lambda_1|^2 \sin 2\alpha. \quad (35)$$

where α is the relative phase of λ_1 and λ_2 . Thus very reasonable choices of the parameters, such as $M_1/M_2 \sim |\lambda_1| \sim \alpha \sim 10^{-1}$, lead to consistency with Eq. (34). We note that since we are considering a $U(1)_x$ gauge symmetry, the Majorana mass terms for

the dark particles are forbidden. Thus, the dark matter asymmetry generated in the early universe would not be washed out by oscillations. The dissipation of the symmetric component of dark matter can be achieved by gauge kinetic energy mixing [22] of $U(1)_x$ and $U(1)_Y$ and via Stueckelberg mass mixing [23–25]. Thus dissipation of the thermally produced $X, X', \tilde{X}, \tilde{X}'$ and their anti-particles occurs from their annihilation via the Z'_x boson exchange coupled with a Breit-Wigner pole enhancement [7, 26–29]. We are able to deplete sufficient amounts of the symmetric component of dark matter (so it is less than 10% of the total dark matter relic density) with a mixing between $U(1)_x$ and $U(1)_Y$ as low as $\delta \sim 0.001$ [7] (where δ is the mixing angle) in the desired mass region of dark matter. Finally, we note that the $U(1)_x$ gaugino λ_x is given a soft mass $\mathcal{L}_{\lambda_x} = m_\lambda \bar{\lambda}_x \lambda_x$. It can then decay into $X\tilde{X}$ or $X'\tilde{X}'$ via the supersymmetric interaction $\mathcal{L} \sim \lambda_x X\tilde{X} + \lambda_x X'\tilde{X}' + h.c.$, where we assume $m_\lambda > m_X + m_{\tilde{X}}$. Thus the gaugino λ_x decays into dark particles and is removed from the low energy spectrum.

Conclusion: In this work we have discussed the possibility that the decay of the inflaton field creates asymmetric dark matter, and the lepton and baryon excess arise as a consequence of transmutation of the asymmetric dark matter. The symmetric component of dark matter is depleted via kinetic mixing between $U(1)_x$ and the hypercharge gauge group and hence annihilates to the SM particles. The Majorana mass terms for the dark particles are forbidden since they carry a $U(1)_x$ gauge charge and thus the asymmetric dark matter generated in the early universe would not be washed out by oscillations and thus sources leptogenesis and baryogenesis.

Acknowledgments: The work of PN is supported in part by the U.S. National Science Foundation (NSF) grants PHY-0757959 and PHY-070467. WZF is supported by funds from The Hong Kong University of Science and Technology. AM is supported by the STFC grant ST/J000418/1. WZF and AM would like to thank Northeastern University for hospitality, and WZF is thankful to HaiPeng for helpful discussions.

-
- [1] A. H. Guth, Phys. Rev. D **23**, 347 (1981).
 - [2] A. D. Linde, Phys. Lett. B **129**, 177 (1983).
 - [3] A. D. Sakharov, Pisma Zh. Eksp. Teor. Fiz. **5**, 32 (1967) [JETP Lett. **5**, 24 (1967)] [Sov. Phys. Usp. **34**, 392 (1991)] [Usp. Fiz. Nauk **161**, 61 (1991)].
 - [4] P. Nath and P. Fileviez Perez, Phys. Rept. **441**, 191 (2007) [hep-ph/0601023].
 - [5] T. R. Dulaney, P. Fileviez Perez and M. B. Wise, Phys.

- Rev. D **83**, 023520 (2011) [arXiv:1005.0617 [hep-ph]].
- [6] M. R. Buckley and L. Randall, JHEP **1109**, 009 (2011) [arXiv:1009.0270 [hep-ph]].
- [7] W. -Z. Feng, P. Nath and G. Peim, Phys. Rev. D **85**, 115016 (2012) [arXiv:1204.5752 [hep-ph]].
- [8] K. Enqvist and A. Mazumdar, Phys. Rept. **380**, 99 (2003) [hep-ph/0209244].
- [9] P. Binetruy and G. R. Dvali, Phys. Lett. B **388**, 241 (1996) [hep-ph/9606342].
- [10] B. Kors and P. Nath, Nucl. Phys. B **711**, 112 (2005) [hep-th/0411201].
- [11] A. Mazumdar and J. Rocher, Phys. Rept. **497**, 85 (2011) [arXiv:1001.0993 [hep-ph]].
- [12] M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986).
- [13] H. Murayama, H. Suzuki, T. Yanagida and J. 'i. Yokoyama, Phys. Rev. Lett. **70**, 1912 (1993).
- [14] M. Plumacher, Z. Phys. C **74**, 549 (1997) [hep-ph/9604229].
- [15] L. Covi, E. Roulet and F. Vissani, Phys. Lett. B **384**, 169 (1996) [hep-ph/9605319].
- [16] W.-Z. Feng and P. Nath, In preparation.
- [17] A. Pilaftsis and T. E. J. Underwood, Nucl. Phys. B **692**, 303 (2004) [hep-ph/0309342].
- [18] W. Buchmuller, P. Di Bari and M. Plumacher, Nucl. Phys. B **643**, 367 (2002); Nucl. Phys. B **665**, 445 (2003).
- [19] D. E. Kaplan, M. A. Luty and K. M. Zurek, Phys. Rev. D **79**, 115016 (2009) [arXiv:0901.4117 [hep-ph]]; M. L. Graesser, I. M. Shoemaker and L. Vecchi, JHEP **1110**, 110 (2011) [arXiv:1103.2771 [hep-ph]].
- [20] J. A. Harvey and M. S. Turner, Phys. Rev. D **42**, 3344 (1990).
- [21] J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).
- [22] B. Holdom, Phys. Lett. B **166**, 196 (1986); Phys. Lett. B **259**, 329 (1991).
- [23] B. Kors and P. Nath, Phys. Lett. B **586**, 366 (2004); JHEP **0412**, 005 (2004); hep-ph/0411406; JHEP **0507**, 069 (2005).
- [24] K. Cheung and T. -C. Yuan, JHEP **0703**, 120 (2007) [hep-ph/0701107].
- [25] D. Feldman, Z. Liu and P. Nath, Phys. Rev. D **75**, 115001 (2007) [hep-ph/0702123 [HEP-PH]].
- [26] K. Griest and D. Seckel, Phys. Rev. D **43**, 3191 (1991).
- [27] P. Gondolo and G. Gelmini, Nucl. Phys. B **360**, 145 (1991).
- [28] R. L. Arnowitt and P. Nath, Phys. Lett. B **299**, 58 (1993); Phys. Rev. Lett. **70**, 3696 (1993).
- [29] D. Feldman, Z. Liu and P. Nath, Phys. Rev. D **79**, 063509 (2009) [arXiv:0810.5762 [hep-ph]].
- [30] In MSSM there are many flat directions which are possible candidates for inflation. For a review see [8].
- [31] In the model only the mass term of the $\hat{N}_i$ superfields ($W \sim \frac{1}{2} M_i \hat{N}_i \hat{N}_i$) violates the lepton number.
- [32] The analysis assumes massless particles running in the loops aside from the heavy field $\hat{N}_2$. Complete results will be presented elsewhere [16].
- [33] The assumption of near degeneracy of the Majorana masses leads to the so-called resonant leptogenesis [17]. However, in the analysis here we do not make use of this mechanism.